



(Patient Sticker)



PHYSICIAN ORDER SET :
OCRELIZUMAB LOAD DOSE (SCHEDULE WEEKS 0, 2)

CDH 208-224 – Approved - Page 1 of 4

Patient: _____ DOB: _____ Gender: _____

Patient Phone #: _____ Height: _____ Weight: _____

Diagnosis: _____ ICD-10 Code: _____

Treatment Start Date: _____

Provider Facility Name: _____ Provider Facility Address: _____

Ordering Provider: _____ Date: _____

Signature: _____

Complete, Sign, and fax this document to: **CDH Central Scheduling at 413-582-2183**

Please include H&P/current medications list/allergies, and ensure that med authorizations have been obtained

Pre-Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 650 mg 650 mg, Oral, Once, Starting S, For 1 Doses	Every 2 weeks		Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) IV 50 mg 50 mg, Intravenous, Once, Starting S, For 1 Doses	Every 2 weeks		Until discont'd
<input type="checkbox"/> methylprednisolone sodium succinate (SOLU-Medrol) IV 100 mg 100 mg, Intravenous, Once, Starting S, For 1 Doses	Every 2 weeks		Until discont'd
<input type="checkbox"/> famotidine (PEPCID) injection 20 mg 20 mg, Intravenous, Once, Starting S, For 1 Doses	Every 2 weeks		Until discont'd

Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> ocrelizumab (OCREVUS) 300 mg in sodium chloride 0.9% 250 mL IVPB 300 mg, Intravenous, Once, Starting H, For 1 Doses <i>Initial infusion: start rate of 30 mL/hr; if there is no reaction, increase the rate by 30 mL/hr increments every 30 minutes to a max rate of 180 mL/hr.</i>	Every 2 weeks		Until discont'd



PHYSICIAN ORDER SET :

OCRELIZUMAB LOAD DOSE (SCHEDULE WEEKS 0, 2)

CDH 208-224 – Approved - Page 2 of 4

Labs

	Interval	Defer Until	Duration
<input type="checkbox"/> CBC and differential Routine, Once, Starting S For 1 Occurrences, If deemed necessary based upon the results of the automated differential, a manual differential may be performed	Every 2 weeks		Until discont'd
<input type="checkbox"/> CD19 Routine, Once, Starting S For 1 Occurrences	Every 2 weeks		Until discont'd
<input type="checkbox"/> MISCELLANEOUS LAB TEST Routine, Once, Starting S For 1 Occurrences <i>MGH - CD19 count, flow cytometry, B cell, NOTE: ONLY lab orders should be entered here. Most lab tests performed in-house and at sendout labs can be found via searching the Facility list. All miscellaneous test requests will be monitored. If known, please enter the tube type as well as the tests requested.</i>	Every 2 weeks		Until discont'd
<input type="checkbox"/> CD20 Routine, Once, Starting S For 1 Occurrences	Every 2 weeks		Until discont'd
<input type="checkbox"/> SPEP panel Routine, Once, Starting S For 1 Occurrences	Every 2 weeks		Until discont'd
<input type="checkbox"/> Comprehensive metabolic panel(BWH , BWF , DFCI , NWH , MEEI , NSMC , SCC , SRH , SHC , NCH , MVH) Routine, Once, Starting S For 1 Occurrences, consists of the following tests: Na, K, Cl, CO2, BUN, Glucose, Creatinine, Calcium, Albumin, Alkaline Phosphatase, Total Bilirubin, Total Protein, ALT, and AST	Every 2 weeks		Until discont'd
<input type="checkbox"/> JC virus antibody inhibition Routine, Once, Starting S For 1 Occurrences	Every 2 weeks		Until discont'd
<input type="checkbox"/> Immunoglobulins IgG, IgA, IgM Routine, Once, Starting S For 1 Occurrences	Every 2 weeks		Until discont'd

Catheter management

	Interval	Defer Until	Duration
<input type="checkbox"/> Line Access Routine, Once, Starting S For 1 Occurrences, As needed. Starting when released. Until Specified. <i>Insert peripheral IV, or access peripheral, or central venous access device, to provide treatment.</i>	PRN		Until discont'd
<input type="checkbox"/> alteplase (CATHFLO) 1 mg/mL injection 2 mg 2 mg, Intracatheter, As needed, line care, For central venous access device requiring clearance. Administer per institutional guidelines. May repeat once per lumen, Starting S	PRN		Until discont'd
<input type="checkbox"/> lidocaine-prilocaine (EMLA) cream Topical, As needed, pre procedure/treatment, Apply prior to the PIV insertion or port access, Starting S	PRN		Until discont'd
<input type="checkbox"/> heparin 100 units/mL flush 5 mL 5 mL, Intravenous, As needed, line care, Line care per institutional guidelines, Starting S	PRN		Until discont'd
<input type="checkbox"/> heparin 10 units/mL flush 3 mL 3 mL, Intravenous, As needed, line care, line care per institutional guidelines, Starting S	PRN		Until discont'd



PHYSICIAN ORDER SET :

OCRELIZUMAB LOAD DOSE (SCHEDULE WEEKS 0, 2)

CDH 208-224 – Approved - Page 3 of 4

Catheter management (continued)

	Interval	Defer Until	Duration
<input type="checkbox"/> heparin 10 units/mL flush 5 mL 5 mL, Intravenous, As needed, line care, line care per institutional guidelines, Starting S	PRN		Until discont'd
<input type="checkbox"/> heparin 1000 units/mL flush 2 mL 2 mL, Intracatheter, As needed, line care, APHERESIS LINE CARE ONLY per institutional guidelines. HEPARIN MUST BE WITHDRAWN FROM EACH LUMEN PRIOR TO FLUSHING OR INFUSING THROUGH THE APHERESIS CATHETER, Starting S	PRN		Until discont'd
<input type="checkbox"/> sodium chloride (NS) 0.9 % syringe flush 3 mL 3 mL, Intravenous, As needed, line care, line care per institutional guidelines, Starting S	PRN		Until discont'd
<input type="checkbox"/> sodium chloride (NS) 0.9 % syringe flush 10 mL 10 mL, Intravenous, As needed, line care, line care per institutional guidelines, Starting S	PRN		Until discont'd
<input type="checkbox"/> sodium chloride (NS) 0.9 % syringe flush 20 mL 20 mL, Intravenous, As needed, line care, line care per institutional guidelines, Starting S	PRN		Until discont'd
<input type="checkbox"/> sodium chloride 0.9% infusion 20 mL/hr, Intravenous, Continuous PRN, other (free text field), Keep vein open to provide treatment, Starting S	PRN		Until discont'd
<input type="checkbox"/> D5W infusion 20 mL/hr, Intravenous, Continuous PRN, other (free text field), Keep vein open to provide treatment, Starting S	PRN		Until discont'd

Emergency Medications

	Interval	Defer Until	Duration
<input type="checkbox"/> Provider and Nurse Communication Routine, Until discontinued, Starting S, Treatment of SEVERE reaction (ANAPHYLAXIS): hypotension, throat swelling, wheezing, respiratory distress, or decreased oxygen saturation. Stop the infusion and treat with Epinephrine FIRST. Notify provider and emergency personnel, administer oxygen as needed, monitor vital signs and proceed with administering adjunct HYPERSENSITIVITY medications as clinically indicated.	PRN		Until discont'd
<input type="checkbox"/> EPINEPHRine injection 0.3 mg 0.3 mg, Intramuscular, As needed, anaphylaxis, Administer FIRST for anaphylaxis. May repeat times 1 dose, Starting S <i>For 2 doses. Pharmacy's Suggested Dose Instructions; Epinephrine 1:1000 is equivalent to 1 mg/mL</i>	PRN		Until discont'd
<input type="checkbox"/> sodium chloride 0.9% bolus 1,000 mL 1,000 mL, Intravenous, Once as needed, other (free text field), For hypotension, Starting S, For 1 Doses	PRN		Until discont'd
<input type="checkbox"/> Oxygen Therapy - Non-Rebreather Routine <i>Select a Mode of Therapy: Non-Rebreather</i>	PRN		Until discont'd
<input type="checkbox"/> Tryptase STAT, Once, Starting S For 1 Occurrences <i>Collect for mild-moderate, or SEVERE reaction</i>	PRN		Until discont'd

Hypersensitivity

	Interval	Defer Until	Duration
<input type="checkbox"/> Provider and Nurse Communication Routine, Until discontinued, Starting S For Until specified, Treatment for mild-moderate infusion reaction: Stop the infusion, notify provider and emergency personnel, administer oxygen as needed, monitor vital signs and proceed with administering medications as clinically indicated. If ANAPHYLAXIS reaction, refer to Emergency Medications section.	PRN		Until discont'd
<input type="checkbox"/> albuterol (ACCUNEB) nebulizer solution 2.5 mg 2.5 mg, Nebulization, Once as needed, shortness of breath, wheezing, wheezing, shortness of breath, Starting S, For 1 Doses	PRN		Until discont'd
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 975 mg 975 mg, Oral, Once as needed, fever, Starting S, For 1 Doses	PRN		Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg	PRN		Until discont'd



PHYSICIAN ORDER SET :

OCRELIZUMAB LOAD DOSE (SCHEDULE WEEKS 0, 2)

CDH 208-224 – Approved - Page 4 of 4

Hypersensitivity (continued)

	Interval	Defer Until	Duration
25 mg, Intravenous, As needed, itching, itching, hives. Begin with 25 mg. If patient has continued reaction, administer additional 25 mg, Starting S			
<input type="checkbox"/> famotidine (PEPCID) injection 20 mg 20 mg, Intravenous, Once as needed, other (free text field), Adjunct treatment for mild-moderate, or SEVERE reaction Hold if: given as premed, Starting S, For 1 Doses	PRN		Until discont'd
<input type="checkbox"/> cetirizine (ZyrTEC) tablet 10-20 mg 10-20 mg, Oral, Once as needed, allergies, Adjunct treatment for mild-moderate, or SEVERE reaction. HOLD IF: giving fexofenadine, Starting S, For 1 Doses	PRN		Until discont'd
<input type="checkbox"/> fexofenadine (ALLEGRA) tablet 90-180 mg 90-180 mg, Oral, Once as needed, allergies, Adjunct treatment for mild-moderate, or SEVERE reaction. HOLD IF: giving cetirizine, Starting S, For 1 Doses	PRN		Until discont'd
<input type="checkbox"/> methylprednisolone sodium succinate (SOLU-Medrol) IV 40 mg 40 mg, Intravenous, Once as needed, other (free text field), Adjunct treatment for mild-moderate, or SEVERE reaction, Starting S, For 1 Doses	PRN		Until discont'd
<input type="checkbox"/> ondansetron (ZOFRAN) injection 4 mg 4 mg, Intravenous, As needed, nausea, vomiting, may repeat x 1 dose, Starting S, For 2 Doses	PRN		Until discont'd
<input type="checkbox"/> meperidine (DEMEROL) injection 25 mg 25 mg, Intravenous, Once as needed, rigors, Starting S, For 1 Doses	PRN		Until discont'd
<input type="checkbox"/> ibuprofen (ADVIL,MOTRIN) tablet 400 mg 400 mg, Oral, Once as needed, mild pain, For 1 Doses	PRN		Until discont'd