



PHYSICIAN ORDER SET :

**BEZLOTOXUMAB**

CDH 208-252 – Approved - Page 1 of 3

Patient: \_\_\_\_\_ DOB: \_\_\_\_\_ Gender: \_\_\_\_\_

Patient Phone #: \_\_\_\_\_ Height: \_\_\_\_\_ Weight: \_\_\_\_\_

Diagnosis: \_\_\_\_\_ ICD-10 Code: \_\_\_\_\_

Treatment Start Date: \_\_\_\_\_

Provider Facility Name: \_\_\_\_\_ Provider Facility Address: \_\_\_\_\_

Ordering Provider: \_\_\_\_\_ Date: \_\_\_\_\_

Signature: \_\_\_\_\_

**Pre-Medications**

	Dose	Interval	Defer Until	Duration
<input type="checkbox"/> <b>acetaminophen (TYLENOL) tablet 650 mg</b> 650 mg, Oral, Once <i>Administer at least 30 minutes prior to principal medication.</i>	650 mg	Once		1 treatment
<input type="checkbox"/> <b>diphenhydrAMINE (BENADRYL) oral 25 mg</b> 25 mg, Oral, Once <i>HOLD IF: given IV. Administer at least 30 minutes prior to principal medication.</i>	25 mg	Once		1 treatment
<input type="checkbox"/> <b>diphenhydrAMINE (BENADRYL) IV 25 mg</b> 25 mg, Intravenous, Once as needed, itching, Patient unable to tolerate PO. <i>HOLD IF: given PO. Administer at least 30 minutes prior to principal medication.</i>	25 mg	Once		1 treatment

**Medications**

	Dose	Interval	Defer Until	Duration
<input type="checkbox"/> <b>bezlotoxumab (ZINPLAVA) 10 mg/kg in sodium chloride 0.9% infusion</b> 10 mg/kg, Intravenous, for 60 Minutes, Once <i>Administer using sterile, nonpyrogenic, low-protein binding 0.2 to 5 micron in-line or add-on filter.</i>	10 mg/kg	Once		1 treatment

**Catheter management**

	Dose	Interval	Defer Until	Duration
<input type="checkbox"/> <b>Line Access</b> Routine <i>As needed. Starting when released. Until Specified.</i>		PRN		Until discont'd
<input type="checkbox"/> <b>alteplase (CATHFLO) 1 mg/mL injection 2 mg</b> 2 mg, Intracatheter, As needed, line care, For central venous access device requiring clearance. Administer per institutional guidelines. May repeat once per lumen.	2 mg	PRN		Until discont'd
<input type="checkbox"/> <b>lidocaine-prilocaine (EMLA) cream</b> Topical, As needed, pre procedure/treatment, Apply prior to the PIV insertion or port access		PRN		Until discont'd



PHYSICIAN ORDER SET :

**BEZLOTOXUMAB**

CDH 208-252 – Approved - Page 2 of 3

**Catheter management (continued)**

	Dose	Interval	Defer Until	Duration
<input type="checkbox"/> <b>heparin 100 units/mL flush 5 mL</b> 5 mL, Intravenous, As needed, line care, Line care per institutional guidelines	5 mL	PRN		Until discont'd
<input type="checkbox"/> <b>heparin 10 units/mL flush 3 mL</b> 3 mL, Intravenous, As needed, line care, Line care per institutional guidelines	3 mL	PRN		Until discont'd
<input type="checkbox"/> <b>heparin 10 units/mL flush 5 mL</b> 5 mL, Intravenous, As needed, line care, Line care per institutional guidelines	5 mL	PRN		Until discont'd
<input type="checkbox"/> <b>heparin 1000 units/mL flush 2 mL</b> 2 mL, Intracatheter, As needed, line care, APHERESIS LINE CARE ONLY per institutional policy. HEPARIN MUST BE WITHDRAWN FROM EACH LUMEN PRIOR TO FLUSHING OR INFUSING THROUGH THE APHERESIS CATHETER	2 mL	PRN		Until discont'd
<input type="checkbox"/> <b>sodium chloride (NS) 0.9 % syringe flush 3 mL</b> 3 mL, Intravenous, As needed, line care, Line care per institutional guidelines	3 mL	PRN		Until discont'd
<input type="checkbox"/> <b>sodium chloride (NS) 0.9 % syringe flush 10 mL</b> 10 mL, Intravenous, As needed, line care, Line care per institutional guidelines	10 mL	PRN		Until discont'd
<input type="checkbox"/> <b>sodium chloride (NS) 0.9 % syringe flush 20 mL</b> 20 mL, Intravenous, As needed, line care, Line care per institutional guidelines	20 mL	PRN		Until discont'd
<input type="checkbox"/> <b>sodium chloride 0.9% infusion</b> 20 mL/hr, Intravenous, Continuous PRN, other (free text field), Keep vein open to provide treatment	20 mL/hr	PRN		Until discont'd
<input type="checkbox"/> <b>D5W infusion</b> 20 mL/hr, Intravenous, Continuous PRN, other (free text field), Keep vein open to provide treatment	20 mL/hr	PRN		Until discont'd

**Emergency Medications/Anaphylaxis**

	Dose	Interval	Defer Until	Duration
<input type="checkbox"/> <b>Provider and Nurse Communication</b> Routine <i>Treatment of SEVERE reaction (ANAPHYLAXIS): hypotension, throat swelling, wheezing, respiratory distress, or decreased oxygen saturation. Stop the infusion and treat with epinephrine FIRST. Notify provider and emergency personnel, administer oxygen as needed, monitor vital signs and proceed with administering adjunct HYPERSENSITIVITY medications as clinically indicated.</i>		PRN		Until discont'd
<input type="checkbox"/> <b>EPINEPHRine injection 0.3 mg</b> 0.3 mg, Intramuscular, As needed, anaphylaxis, Administer FIRST for anaphylaxis. May repeat times 1 dose <i>For 2 doses. Pharmacy's Suggested Dose Instructions; Epinephrine 1:1000 is equivalent to 1 mg/mL</i>	.3 mg	PRN		Until discont'd
<input type="checkbox"/> <b>sodium chloride 0.9% bolus 1,000 mL</b> 1,000 mL, Intravenous, Once as needed, other (free text field), For hypotension	1000 mL	PRN		Until discont'd
<input type="checkbox"/> <b>Oxygen Therapy - Non-Rebreather</b> Routine <i>Select a Mode of Therapy: Non-Rebreather</i>		PRN		Until discont'd

**Hypersensitivity**

	Dose	Interval	Defer Until	Duration
<input type="checkbox"/> <b>Provider and Nurse Communication</b> Routine <i>Treatment for mild-moderate infusion reaction: Stop the infusion, notify provider and emergency personnel, administer oxygen as needed, monitor vital signs and proceed with administering medications as clinically indicated. If ANAPHYLAXIS reaction, refer to Emergency Medications section.</i>		PRN		Until discont'd
<input type="checkbox"/> <b>albuterol (ACCUNEB) nebulizer solution 2.5 mg</b> 2.5 mg, Nebulization, Once as needed, shortness of breath/dyspnea, wheezing, wheezing, shortness of breath	2.5 mg	PRN		Until discont'd



PHYSICIAN ORDER SET :

**BEZLOTOXUMAB**

CDH 208-252 – Approved - Page 3 of 3

**Hypersensitivity (continued)**

	Dose	Interval	Defer Until	Duration
<input type="checkbox"/> <b>acetaminophen (TYLENOL) tablet 975 mg</b> 975 mg, Oral, Once as needed, fever	975 mg	PRN		Until discont'd
<input type="checkbox"/> <b>diphenhydrAMINE (BENADRYL) injection 25 mg</b> 25 mg, Intravenous, As needed, itching, itching, hives. Begin with 25 mg. If patient has continued reaction, administer additional 25 mg	25 mg	PRN		Until discont'd
<input type="checkbox"/> <b>famotidine (PEPCID) injection 20 mg</b> 20 mg, Intravenous, Once as needed, other (free text field), Adjunct treatment for mild-moderate, or SEVERE reaction Hold if: given as premed	20 mg	PRN		Until discont'd
<input type="checkbox"/> <b>cetirizine (ZyrTEC) tablet 10 mg</b> 10 mg, Oral, Once as needed, allergies, Adjunct treatment for mild-moderate, or SEVERE reaction <i>HOLD IF: given fexofenadine.</i>	10 mg	PRN		Until discont'd
<input type="checkbox"/> <b>fexofenadine (ALLEGRA) tablet 180 mg</b> 180 mg, Oral, Once as needed, allergies, Adjunct treatment for mild-moderate, or SEVERE reaction <i>HOLD IF: given cetirizine.</i>	180 mg	PRN		Until discont'd
<input type="checkbox"/> <b>methylprednisolone sodium succinate (SOLU-Medrol) IV 40 mg</b> 40 mg, Intravenous, Once as needed, other (free text field), Adjunct treatment for mild-moderate, or SEVERE reaction	40 mg	PRN		Until discont'd
<input type="checkbox"/> <b>ondansetron (ZOFRAN) injection 4 mg</b> 4 mg, Intravenous, As needed, nausea, vomiting, may repeat x 1 dose	4 mg	PRN		Until discont'd