



(Patient Sticker)



PHYSICIAN ORDER SET :

BELATACEPT LOAD DOSE

CDH 208-256 – Approved - Page 1 of 4

Patient: _____

DOB: _____

Gender: _____

Patient Phone #: _____

Height: _____

Weight: _____

Diagnosis: _____

ICD-10 Code: _____

Treatment Start Date: _____

Provider Facility Name: _____

Provider Facility Address: _____

Ordering Provider: _____

Date: _____

Signature: _____

Hydrations

	Dose	Interval	Defer Until	Duration
<input type="checkbox"/> lactated Ringers infusion 250 mL/hr, Intravenous, Continuous	250 mL/hr	Every 2 weeks		5 treatments
<input type="checkbox"/> D5-NS infusion 250 mL/hr, Intravenous, Continuous	250 mL/hr	Every 2 weeks		5 treatments
<input type="checkbox"/> D5-1/2 NS infusion 250 mL/hr, Intravenous, Continuous	250 mL/hr	Every 2 weeks		5 treatments
<input type="checkbox"/> D5W infusion 250 mL/hr, Intravenous, Continuous	250 mL/hr	Every 2 weeks		5 treatments
<input type="checkbox"/> sodium chloride 0.9% infusion 250 mL/hr, Intravenous, Continuous	250 mL/hr	Every 2 weeks		5 treatments
<input type="checkbox"/> sodium chloride 0.9% bolus 500 mL 500 mL, Intravenous, for 30 Minutes, Continuous	500 mL	Every 2 weeks		5 treatments
<input type="checkbox"/> sodium chloride 0.9% bolus 1,000 mL 1,000 mL, Intravenous, for 60 Minutes, Continuous	1000 mL	Every 2 weeks		5 treatments



PHYSICIAN ORDER SET :

BELATACEPT LOAD DOSE

CDH 208-256 – Approved - Page 2 of 4

Labs Medications

	Dose	Interval	Defer Until	Duration
<input type="checkbox"/> belatacept (NULOJIX) IV 10 mg/kg	10 mg/kg	Every 2 weeks		5 treatments
10 mg/kg, Intravenous, Continuous Follow your institutional guidelines.				

Labs

	Dose	Interval	Defer Until	Duration
<input type="checkbox"/> Comprehensive metabolic panel		Every 2 weeks		Until discont'd
Routine, consists of the following tests: Na, K, Cl, CO2, BUN, Glucose, Creatinine, Calcium, Albumin, Alkaline Phosphatase, Total Bilirubin, Total Protein, ALT, and AST				
<input type="checkbox"/> CBC and differential		Every 2 weeks		Until discont'd
Routine, If deemed necessary based upon the results of the automated differential, a manual differential may be performed				
<input type="checkbox"/> Urinalysis		Every 2 weeks		Until discont'd
Except NWH Urinalysis sediments are performed on all urines that have positive findings (blood, WBC, or protein) on the chemical urinalysis. Urine sediment does not need to be ordered separately.				

Catheter management

	Dose	Interval	Defer Until	Duration
<input type="checkbox"/> Line Access		PRN		Until discont'd
Routine As needed. Until Specified. Insert peripheral IV, or access peripheral, or central venous access device, to provide treatment.				
<input type="checkbox"/> alteplase (CATHFLO) 1 mg/mL injection 2 mg	2 mg	PRN		Until discont'd
2 mg, Intracatheter, As needed, line care For central venous access device requiring clearance. Administer per institutional guidelines. May repeat once per lumen.				
<input type="checkbox"/> lidocaine-prilocaine (EMLA) cream	___	PRN		Until discont'd
Topical, As needed, pre procedure/treatment, apply prior to the PIV insertion or port access				
<input type="checkbox"/> heparin 100 units/mL flush 5 mL	5 mL	PRN		Until discont'd
5 mL, Intravenous, As needed, line care, Line care per institutional guidelines				
<input type="checkbox"/> heparin 10 units/mL flush 3 mL	3 mL	PRN		Until discont'd
3 mL, Intravenous, As needed, line care, Line care per institutional guidelines				
<input type="checkbox"/> heparin 10 units/mL flush 5 mL	5 mL	PRN		Until discont'd
5 mL, Intravenous, As needed, line care, Line care per institutional guidelines				
<input type="checkbox"/> heparin 1000 units/mL flush 2 mL	2 mL	PRN		Until discont'd
2 mL, Intracatheter, As needed, line care, APHERESIS LINE CARE ONLY per institutional policy. HEPARIN MUST BE WITHDRAWN FROM EACH LUMEN PRIOR TO FLUSHING OR INFUSING THROUGH THE APHERESIS CATHETER				
<input type="checkbox"/> sodium chloride (NS) 0.9 % syringe flush 3 mL	3 mL	PRN		Until discont'd
3 mL, Intravenous, As needed, line care, Line care per institutional guidelines				



PHYSICIAN ORDER SET :

BELATACEPT LOAD DOSE

CDH 208-256 – Approved - Page 3 of 4

Catheter management (continued)

	Dose	Interval	Defer Until	Duration
<input type="checkbox"/> sodium chloride (NS) 0.9 % syringe flush 10 mL 10 mL, Intravenous, As needed, line care, Line care per institutional guidelines	10 mL	PRN		Until discont'd
<input type="checkbox"/> sodium chloride (NS) 0.9 % syringe flush 20 mL 20 mL, Intravenous, As needed, line care, Line care per institutional guidelines	20 mL	PRN		Until discont'd
<input type="checkbox"/> sodium chloride 0.9% infusion 20 mL/hr, Intravenous, Continuous PRN, other (free text field), Keep vein open to provide treatment	20 mL/hr	PRN		Until discont'd
<input type="checkbox"/> D5W infusion 20 mL/hr, Intravenous, Continuous PRN, other (free text field), Keep vein open to provide treatment	20 mL/hr	PRN		Until discont'd

Emergency Medications/Anaphylaxis

	Dose	Interval	Defer Until	Duration
<input type="checkbox"/> Provider and Nurse Communication Routine <i>Treatment of SEVERE reaction (ANAPHYLAXIS): hypotension, throat swelling, wheezing, respiratory distress, or decreased oxygen saturation. Stop the infusion and treat with epinephrine FIRST. Notify provider and emergency personnel, administer oxygen as needed, monitor vital signs and proceed with administering adjunct HYPERSENSITIVITY medications as clinically indicated.</i>		PRN		Until discont'd
<input type="checkbox"/> EPINEPHrine injection 0.3 mg 0.3 mg, Intramuscular, As needed, anaphylaxis, Administer FIRST for anaphylaxis. May repeat times 1 dose <i>For 2 doses. Pharmacy's Suggested Dose Instructions; Epinephrine 1:1000 is equivalent to 1 mg/mL</i>	.3 mg	PRN		Until discont'd
<input type="checkbox"/> sodium chloride 0.9% bolus 1,000 mL 1,000 mL, Intravenous, Once as needed, other (free text field), Hypotension	1000 mL	PRN		Until discont'd
<input type="checkbox"/> Oxygen Therapy - Non-Rebreather Routine <i>Select a Mode of Therapy: Non-Rebreather</i>		PRN		Until discont'd

Hypersensitivity

	Dose	Interval	Defer Until	Duration
<input type="checkbox"/> Provider and Nurse Communication Routine <i>Treatment for mild-moderate infusion reaction: Stop the infusion, notify provider and emergency personnel, administer oxygen as needed, monitor vital signs and proceed with administering medications as clinically indicated. If ANAPHYLAXIS reaction, refer to Emergency Medications section.</i>		PRN		Until discont'd
<input type="checkbox"/> albuterol (ACCUNEB) nebulizer solution 2.5 mg 2.5 mg, Nebulization, Once as needed, wheezing, shortness of breath	2.5 mg	PRN		Until discont'd
<input type="checkbox"/> acetaminophen (TYLENOL) tablet 975 mg 975 mg, Oral, Once as needed, fever	975 mg	PRN		Until discont'd
<input type="checkbox"/> diphenhydrAMINE (BENADRYL) injection 25 mg 25 mg, Intravenous, As needed, itching, itching, hives. Begin with 25 mg. If patient has continued reaction, administer additional 25 mg	25 mg	PRN		Until discont'd
<input type="checkbox"/> famotidine (PEPCID) injection 20 mg 20 mg, Intravenous, Once as needed, other (free text field), Adjunct treatment for mild-moderate, or SEVERE reaction Hold if: given as premed	20 mg	PRN		Until discont'd



(Patient Sticker)

PHYSICIAN ORDER SET :

BELATACEPT LOAD DOSE

CDH 208-256 – Approved - Page 4 of 4

Hypersensitivity (continued)

	Dose	Interval	Defer Until	Duration
<input type="checkbox"/> cetirizine (ZyrTEC) tablet 10 mg 10 mg, Oral, Once as needed, other (free text field), Adjunct treatment for mild-moderate, or SEVERE reaction. <i>HOLD IF: given fexofenadine.</i>	10 mg	PRN		Until discont'd
<input type="checkbox"/> fexofenadine (ALLEGRA) tablet 180 mg 180 mg, Oral, Once as needed, allergies, Adjunct treatment for mild-moderate, or SEVERE reaction. <i>HOLD IF: given cetirizine.</i>	180 mg	PRN		Until discont'd
<input type="checkbox"/> methylprednisolone sodium succinate (SOLU-Medrol) IV 40 mg 40 mg, Intravenous, Once as needed, other (free text field), Adjunct treatment for mild-moderate, or SEVERE reaction	40 mg	PRN		Until discont'd
<input type="checkbox"/> ondansetron (ZOFRAN) injection 4 mg 4 mg, Intravenous, As needed, nausea, vomiting, may repeat x 1 dose	4 mg	PRN		Until discont'd